

The type I conjecture on trees

Semal Lancelot

UCLouvain-FNRS GRANT

October 7, 2021

Unitary representations

Unitary representations

Why do we study unitary representations ?

Unitary representations

Why do we study unitary representations ?

Groups G

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups G

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups $G \Rightarrow$ **Haar** measure μ

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups $G \Rightarrow$ **Haar** measure μ
 \Rightarrow actions on **Hilbert** spaces $G \curvearrowright \mathcal{H}$

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups $G \Rightarrow$ **Haar** measure μ
 \Rightarrow actions on **Hilbert** spaces $G \curvearrowright \mathcal{H}$

The left-regular representation

Let G be a locally compact group we define a group action $G \curvearrowright L^2(G, \mu)$:

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups $G \Rightarrow$ **Haar** measure μ
 \Rightarrow actions on **Hilbert** spaces $G \curvearrowright \mathcal{H}$

The left-regular representation

Let G be a locally compact group we define a group action $G \curvearrowright L^2(G, \mu)$:

$$\lambda_G(g)f(h) = f(g^{-1}h) \quad \forall g, h \in G.$$

Unitary representations

Why do we study unitary representations ?

Groups $G \Rightarrow$ actions $G \curvearrowright X$.

Locally compact groups $G \Rightarrow$ **Haar** measure μ
 \Rightarrow actions on **Hilbert** spaces $G \curvearrowright \mathcal{H}$

The left-regular representation

Let G be a locally compact group we define a group action $G \curvearrowright L^2(G, \mu)$:

$$\lambda_G(g)f(h) = f(g^{-1}h) \quad \forall g, h \in G.$$

This define a continuous **faithfull** action of G on a Hilbert space.

Unitary representations of locally compact groups

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Equivalence representations

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Equivalence representations

Two representations π and σ are said to be **equivalent** when there exists a unitary operator :

$$\mathcal{U} : \mathcal{H}_\pi \rightarrow \mathcal{H}_\sigma$$

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Equivalence representations

Two representations π and σ are said to be **equivalent** when there exists a unitary operator :

$$\mathcal{U} : \mathcal{H}_\pi \rightarrow \mathcal{H}_\sigma$$

such that

$$\mathcal{U}\pi(g) = \sigma(g)\mathcal{U} \quad \forall g \in G.$$

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Irreducible representations

Unitary representations of locally compact groups

Representations

A **unitary representation** of a locally compact group G is a continuous group homomorphism :

$$\pi : G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$$

where \mathcal{H}_π is a complex Hilbert space.

Irreducible representations

A representation is called **irreducible** when the only **closed** invariant subspaces $W \subseteq \mathcal{H}_\pi$ (e.g $\pi(G)W \subseteq W$) are $\{0\}$ and \mathcal{H}_π .

Decomposition of representations

Decomposition of representations

Decomposition of representations

Let G be a finite group.

Decomposition of representations

Decomposition of representations

Let G be a **finite** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Remark

The study of representations of **compact** groups reduces to the study of **irreducible** representations.

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Let G be a **locally compact** group.

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Let G be a **locally compact** group.

Every representation π of G **decomposes (possibly not uniquely)**

Decomposition of representations

Decomposition of representations

Let G be a **compact** group.

Every representation π of G **decomposes uniquely** as a **direct sum** of **irreducible** representations of G :

$$\pi \simeq \bigoplus_{i \in \mathbb{N}} n_i \sigma_i.$$

Let G be a **locally compact** group.

Every representation π of G **decomposes (possibly not uniquely)** as a **direct integral** of **irreducible** representations of G :

$$\pi \simeq \int_{\widehat{G}} \sigma \, d\nu_{\pi}(\sigma).$$

Right building blocks

Right building blocks

Factorial representations

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Examples

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Examples

Let σ be an **irreducible** representation.

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Examples

Let σ be an **irreducible** representation. The representation:

$$\pi \simeq n\sigma$$

is a **factorial** representation for every $n = \infty, 1, 2, \dots$

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Examples

Let F_2 be the free group with two generators.

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Examples

Let F_2 be the free group with two generators. The regular representation

$$(\lambda_{F_2}, L^2(F_2, \mu))$$

is a **factorial** representation.

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Non-example

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Non-example

Let σ and τ be **not equivalent irreducible** representations.

Right building blocks

Factorial representations

A representation is said to be **factorial** when for any decomposition

$$\pi \simeq \sigma_1 \oplus \sigma_2$$

we have that

$$\sigma_1 \leq \infty \sigma_2 \text{ and } \sigma_2 \leq \infty \sigma_1.$$

Non-example

Let σ and τ be **not equivalent irreducible** representations. The representation:

$$\pi \simeq n_1 \sigma \oplus n_2 \tau$$

is a **not factorial** for any $n_1, n_2 \geq 1$.

Decomposition of representations

Decomposition of representations

Decomposition of representations

Let G be a **locally compact** group.

Every representation π of G **decomposes (possibly not uniquely)** as a **direct integral** of **irreducible** representations of G :

$$\pi \simeq \int_{\widehat{G}} \sigma \, d\nu_{\pi}(\sigma).$$

Decomposition of representations

Decomposition of representations

Let G be a **locally compact** group.

Every representation π of G **decomposes (possibly not uniquely)** as a **direct integral** of **irreducible** representations of G :

$$\pi \simeq \int_{\widehat{G}} \sigma \, d\nu_{\pi}(\sigma).$$

Every representation π of G **decomposes uniquely**

Decomposition of representations

Decomposition of representations

Let G be a **locally compact** group.

Every representation π of G **decomposes (possibly not uniquely)** as a **direct integral** of **irreducible** representations of G :

$$\pi \simeq \int_{\widehat{G}} \sigma \, d\nu_{\pi}(\sigma).$$

Every representation π of G **decomposes uniquely** as a **direct integral** of **factorial** representations of G :

$$\pi \simeq \int_{\text{Fact}(G)} \sigma \, d\nu_{\pi}(\sigma).$$

Type I groups

Type I groups

A locally compact group is called a **type I** group when every of its factorial representations is of the form:

$$\pi \simeq n\sigma$$

for some irreducible representation σ and $n = \infty, 1, 2, \dots$

Type I groups

A locally compact group is called a **type I** group when every of its factorial representations is of the form:

$$\pi \simeq n\sigma$$

for some irreducible representation σ and $n = \infty, 1, 2, \dots$

Remark

For **type I** groups

Type I groups

A locally compact group is called a **type I** group when every of its factorial representations is of the form:

$$\pi \simeq n\sigma$$

for some irreducible representation σ and $n = \infty, 1, 2, \dots$

Remark

For **type I** groups the decomposition of representations into **direct integral** of **irreducible** representations is **unique**.

Type I groups

Type I groups

Examples of type I groups

Type I groups

Examples of type I groups

- **Finite** groups.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.
- **Semisimple algebraic** groups over local fields.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.
- **Semisimple algebraic** groups over local fields.
- The full **automorphism** group of a **semiregular tree**.

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.
- **Semisimple algebraic** groups over local fields.
- The full **automorphism** group of a **semiregular tree**.

Examples of non-type I groups

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.
- **Semisimple algebraic** groups over local fields.
- The full **automorphism** group of a **semiregular tree**.

Examples of non-type I groups

- **Free** groups with rank ≥ 2 .

Type I groups

Examples of type I groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Virtually abelian discrete** groups.
- **Semisimple algebraic** groups over local fields.
- The full **automorphism** group of a **semiregular tree**.

Examples of non-type I groups

- **Free** groups with rank ≥ 2 .
- **Non-virtually abelian discrete** groups.

Type I conjecture on trees

Type I conjecture on trees

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I**

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

A closed subgroup $G \leq \text{Aut}(T)$ is **amenable** if and only if G satisfies one of the following :

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

A closed subgroup $G \leq \text{Aut}(T)$ is **amenable** if and only if G satisfies one of the following :

- G fixes a **vertex** of T .

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

A closed subgroup $G \leq \text{Aut}(T)$ is **amenable** if and only if G satisfies one of the following :

- G fixes a **vertex** of T .
- G stabilizes an **edge** of T .

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

A closed subgroup $G \leq \text{Aut}(T)$ is **amenable** if and only if G satisfies one of the following :

- G fixes a **vertex** of T .
- G stabilizes an **edge** of T .
- G fixes a **point in the boundary**.

Type I conjecture on trees

Which closed subgroup of $\text{Aut}(T)$ is **type I** among **non-amenable** groups ?

Amenability criterion in trees

A closed subgroup $G \leq \text{Aut}(T)$ is **amenable** if and only if G satisfies one of the following :

- G fixes a **vertex** of T .
- G stabilizes an **edge** of T .
- G fixes a **point in the boundary**.
- G stabilizes a **pair of points in the boundary**.

The type I conjecture on trees

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a closed non-amenable subgroup acting minimally on T .

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a closed non-amenable subgroup acting minimally on T .

Then, G acts **transitively** on the boundary if and only if it is **type I**.

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a closed non-amenable subgroup acting minimally on T .

Then, G acts **transitively** on the boundary if and only if it is **type I**.

Type I conjecture on trees

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a closed non-amenable subgroup acting minimally on T .

Then, G acts **transitively** on the boundary if and only if it is **type I**.

Type I conjecture on trees

Let Δ be a locally finite thick building of type $I_2(\infty)$ and let $G \leq \text{Aut}(\Delta)$ be a closed non-amenable subgroup acting minimally on Δ .

The type I conjecture on trees

Type I conjecture on trees (Nebbia-Houdayer-Raum)

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a closed non-amenable subgroup acting minimally on T .

Then, G acts **transitively** on the boundary if and only if it is **type I**.

Type I conjecture on trees

Let Δ be a locally finite thick building of type $I_2(\infty)$ and let $G \leq \text{Aut}(\Delta)$ be a closed non-amenable subgroup acting minimally on Δ .

Then, G comes from a **B-N** pair if and only if it is **type I**.

Part I : G acts transitively on the boundary then G is type I.

Part I : G acts transitively on the boundary then G is type I.

Partially solved

CCR groups

CCR groups

CCR groups

CCR groups

A locally compact group G is called **CCR**

CCR groups

A locally compact group G is called **CCR** if every **irreducible** representation π of G is **admissible**

CCR groups

A locally compact group G is called **CCR** if every **irreducible** representation π of G is **admissible** i.e. for every compact subgroup $K \leq G$

CCR groups

A locally compact group G is called **CCR** if every **irreducible** representation π of G is **admissible** i.e. for every compact subgroup $K \leq G$ the multiplicities n_i appearing in the decomposition :

$$\pi|_K \simeq \bigoplus_i n_i \sigma_i$$

into irreducibles of K

CCR groups

A locally compact group G is called **CCR** if every **irreducible** representation π of G is **admissible** i.e. for every compact subgroup $K \leq G$ the multiplicities n_i appearing in the decomposition :

$$\pi|_K \simeq \bigoplus_i n_i \sigma_i$$

into irreducibles of K are all **finite**.

CCR groups

CCR groups

A locally compact group G is called **CCR** if every **irreducible** representation π of G is **admissible** i.e. for every compact subgroup $K \leq G$ the multiplicities n_i appearing in the decomposition :

$$\pi|_K \simeq \bigoplus_i n_i \sigma_i$$

into irreducibles of K are all **finite**.

Criterion for “type I”ness

CCR groups are **type I**.

Examples of CCR groups

- **Finite** groups.
- **Compact** groups.
- **Abelian** groups.
- **Semisimple algebraic** groups over local fields.
- The full **automorphism** group of a **semiregular tree**.

Representations of $\text{Aut}(T)$

The topology of $G \leq \text{Aut}(T)$

Representations of $\text{Aut}(T)$

The topology of $G \leq \text{Aut}(T)$

Let $G \leq \text{Aut}(T)$ be a closed subgroup.


Representations of $\text{Aut}(T)$

The topology of $G \leq \text{Aut}(T)$

Let $G \leq \text{Aut}(T)$ be a closed subgroup. The sets

$$\text{Fix}_{\text{Aut}(T)}(\mathcal{T}) = \{g \in \text{Aut}(T) \mid \forall v \in \mathcal{T} : gv = v\}$$

where \mathcal{T} is a complete finite subtree




Representations of $\text{Aut}(T)$

The topology of $G \leq \text{Aut}(T)$

Let $G \leq \text{Aut}(T)$ be a closed subgroup. The sets

$$\text{Fix}_{\text{Aut}(T)}(\mathcal{T}) = \{g \in \text{Aut}(T) \mid \forall v \in \mathcal{T} : gv = v\}$$

where \mathcal{T} is a complete finite subtree form a **b.o.n.c.o.** of G .



The irreducible representations of $\text{Aut}(T)$

The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G .

The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero $\text{Fix}_G(\mathcal{T})$ -invariant vector** $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero** $\text{Fix}_G(\mathcal{T})$ -**invariant vector** $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

Proof.



The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero** $\text{Fix}_G(\mathcal{T})$ -**invariant vector** $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

Proof.

Let $\xi \in \mathcal{H}_\pi$ be non trivial.



The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero** $\text{Fix}_G(\mathcal{T})$ -invariant vector $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

Proof.

Let $\xi \in \mathcal{H}_\pi$ be non trivial. Now, $\int_{\text{Fix}_G(\mathcal{T})} \pi(g)\xi d\mu(g)$ is $\text{Fix}_G(\mathcal{T})$ -invariant.



The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero** $\text{Fix}_G(\mathcal{T})$ -invariant vector $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

Proof.

Let $\xi \in \mathcal{H}_\pi$ be non trivial. Now, $\int_{\text{Fix}_G(\mathcal{T})} \pi(g)\xi d\mu(g)$ is $\text{Fix}_G(\mathcal{T})$ -invariant. The function $g \mapsto \langle \pi(g)\xi, \xi \rangle$ is continuous and non-zero at $g = 1_G$.



The irreducible representations of $\text{Aut}(T)$

Lemma

Let π be a representation of G . Then, there **exists** a complete finite subtree \mathcal{T} of T such that π admits a **non-zero** $\text{Fix}_G(\mathcal{T})$ -invariant vector $\xi \in \mathcal{H}_\pi$ i.e. $\pi(k)\xi = \xi \ \forall k \in \text{Fix}_G(\mathcal{T})$.

Proof.

Let $\xi \in \mathcal{H}_\pi$ be non trivial. Now, $\int_{\text{Fix}_G(\mathcal{T})} \pi(g)\xi d\mu(g)$ is $\text{Fix}_G(\mathcal{T})$ -invariant. The function $g \mapsto \langle \pi(g)\xi, \xi \rangle$ is continuous and non-zero at $g = 1_G$. \exists a big enough \mathcal{T} such that $\int_{\text{Fix}_G(\mathcal{T})} \pi(g)\xi d\mu(g) \neq 0$. □

Minimal subtree

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a vertex

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a vertex π is called **spherical**.

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a vertex π is called **spherical**.
- an edge

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
- an **edge** π is called **special**.

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
- an **edge** π is called **special**.
- **bigger than an edge**

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
- an **edge** π is called **special**.
- **bigger than an edge** π is called **cuspidal**.

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
 G is transitive on the boundary
- an **edge** π is called **special**.
 G is transitively on the boundary
- **bigger than an edge** π is called **cuspidal**.

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
 G is transitive on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- an **edge** π is called **special**.
 G is transitively on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- **bigger than an edge** π is called **cuspidal**.

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
 G is transitive on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- an **edge** π is called **special**.
 G is transitively on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- **bigger than an edge** π is called **cuspidal**.
 G has the Tits independence property (Ol'Shanskii-Amann-Ciobotaru)

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
 G is transitive on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- an **edge** π is called **special**.
 G is transitively on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- **bigger than an edge** π is called **cuspidal**.
 G has the Tits independance property (Ol'Shanskii-Amann-Ciobotaru)
or is a Radu groups (Semal)

Minimal subtree

Given a representation π of G , what is the **smallest** complete finite subtree \mathcal{T} for which it admits invariant vectors ?

Three types

If the minimal subtree \mathcal{T} is :

- a **vertex** π is called **spherical**.
 G is transitive on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- an **edge** π is called **special**.
 G is transitively on the boundary \Rightarrow classified and **CCR**. (Ol'Shanskii)
- **bigger than an edge** π is called **cuspidal**.
 G has the Tits independance property (Ol'Shanskii-Amann-Ciobotaru)
or is a Radu groups (Semal) \Rightarrow classified and **CCR**.

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Remark

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Remark

The set Θ of integers $d \in \mathbb{N}$ for which a 2 transitive subgroup of $\text{Sym}(d)$ contains the alternating group $\text{Alt}(d)$ is

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Remark

The set Θ of integers $d \in \mathbb{N}$ for which a 2 transitive subgroup of $\text{Sym}(d)$ contains the alternating group $\text{Alt}(d)$ is **asymptotically dense** in \mathbb{N} .

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Remark

The set Θ of integers $d \in \mathbb{N}$ for which a 2 transitive subgroup of $\text{Sym}(d)$ contains the alternating group $\text{Alt}(d)$ is **asymptotically dense** in \mathbb{N} .
The 10 smallest element of Θ being 34, 35, 39, 45, 46, 51, 52, 55, 56, 58.

Classification (Radu 2017)

When T is a (d_0, d_1) -semiregular tree with $d_0, d_1 \geq 6$, Radu has completely classified non-compact closed subgroup acting **transitively on the boundary** ∂T and whose **local action** at every vertex contains the **alternating group**.

Remark

The set Θ of integers $d \in \mathbb{N}$ for which a 2 transitive subgroup of $\text{Sym}(d)$ contains the alternating group $\text{Alt}(d)$ is **asymptotically dense** in \mathbb{N} .
The 10 smallest element of Θ being 34, 35, 39, 45, 46, 51, 52, 55, 56, 58.

This **solves the part I of the type I conjecture** for numerous semi-regular trees such as the $(34, 55)$ -semiregular tree.

Part II of the type I conjecture

Part II : G is type I then it acts transitively on the boundary.

Part II of the type I conjecture

Part II : G is type I then it acts transitively on the boundary.

Completely solved !

Part II of the type I conjecture

Part II of the type I conjecture

Theorem (Nebbia 1999)

Part II of the type I conjecture

Theorem (Nebbia 1999)

A **CCR** subgroup of $\text{Aut}(T)$

Part II of the type I conjecture

Theorem (Nebbia 1999)

A **CCR** subgroup of $\text{Aut}(T)$ acts **transitively on the boundary**.

Part II of the type I conjecture

Theorem (Nebbia 1999)

A **CCR** subgroup of $\text{Aut}(T)$ acts **transitively on the boundary**.

Theorem (Houdayer-Raum 2016)

Let $G \leq \text{Aut}(T)$ be a closed non-amenable **type I** subgroup acting minimally on T then it acts

Part II of the type I conjecture

Theorem (Nebbia 1999)

A **CCR** subgroup of $\text{Aut}(T)$ acts **transitively on the boundary**.

Theorem (Houdayer-Raum 2016)

Let $G \leq \text{Aut}(T)$ be a closed non-amenable **type I** subgroup acting minimally on T then it acts **locally 2-transitively**.

Part II of the type I conjecture

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.
- 2 G is **type I**.

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.
- 2 G is **type I**.

Then, G contains a cocompact amenable subgroup.

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.
- 2 G is **type I**.

Then, G contains a cocompact amenable subgroup.

Corollary

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.
- 2 G is **type I**.

Then, G contains a cocompact amenable subgroup.

Corollary

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a **Type I** closed non-amenable subgroup acting minimally on T .

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- ① G contains a **cocompact lattice**.
- ② G is **type I**.

Then, G contains a cocompact amenable subgroup.

Corollary

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a **Type I** closed non-amenable subgroup acting minimally on T . Then, G has a cocompact amenable subgroup

Part II of the type I conjecture

Theorem (Kalantar-Monod-Caprace (Fresh of the week))

Let G be a hyperbolic locally compact group such that:

- 1 G contains a **cocompact lattice**.
- 2 G is **type I**.

Then, G contains a cocompact amenable subgroup.

Corollary

Let T be a locally finite thick tree and let $G \leq \text{Aut}(T)$ be a **Type I** closed non-amenable subgroup acting minimally on T . Then, G has a cocompact amenable subgroup and hence acts transitively on the boundary of T .

Theorem

Let $G \leq \text{Aut}(T)$ be a **Type I** closed non-amenable subgroup acting minimally on T . Then, G acts **transitively** on the boundary of the tree.

Thank you

Thank you !!