

# Divergence in Coxeter groups

Anne Thomas

University of Sydney

Buildings 2021  
Magdeburg

# Outline

This is joint work with Pallavi Dani, Yusra Naqvi, Dibyendu Roy and Ignat Soroko.

# Outline

This is joint work with Pallavi Dani, Yusra Naqvi, Dibyendu Roy and Ignat Soroko.

1. Divergence

# Outline

This is joint work with Pallavi Dani, Yusra Naqvi, Dibyendu Roy and Ignat Soroko.

1. Divergence
2. Right-angled Coxeter groups

# Outline

This is joint work with Pallavi Dani, Yusra Naqvi, Dibyendu Roy and Ignat Soroko.

1. Divergence
2. Right-angled Coxeter groups
3. General Coxeter groups

# Outline

This is joint work with Pallavi Dani, Yusra Naqvi, Dibyendu Roy and Ignat Soroko.

1. Divergence
2. Right-angled Coxeter groups
3. General Coxeter groups (infinite and non-affine)

# Divergence of geodesics

Let  $(X, d)$  be a one-ended geodesic metric space.

## Divergence of geodesics

Let  $(X, d)$  be a one-ended geodesic metric space.

Let  $\gamma_1, \gamma_2 : [0, \infty) \rightarrow X$  be geodesic rays with the same basepoint  $\gamma_1(0) = \gamma_2(0) = x_0$ .



# Divergence of geodesics

Let  $(X, d)$  be a one-ended geodesic metric space.

Let  $\gamma_1, \gamma_2 : [0, \infty) \rightarrow X$  be geodesic rays with the same basepoint  $\gamma_1(0) = \gamma_2(0) = x_0$ .

## Question

*How fast do  $\gamma_1$  and  $\gamma_2$  move away from each other?*

# Divergence of geodesics

Let  $(X, d)$  be a one-ended geodesic metric space.

Let  $\gamma_1, \gamma_2 : [0, \infty) \rightarrow X$  be geodesic rays with the same basepoint  $\gamma_1(0) = \gamma_2(0) = x_0$ .

## Question

*How fast do  $\gamma_1$  and  $\gamma_2$  move away from each other?*

## Definition (Gromov 1980s)


The **divergence** of  $\gamma_1$  and  $\gamma_2$  at time  $r$  is

$$\text{div}(\gamma_1, \gamma_2, r) := \inf_p \text{length}(p)$$

where the infimum is taken over all rectifiable paths  $p$  in  $X \setminus \text{Ball}(x_0, r)$  connecting  $\gamma_1(r)$  and  $\gamma_2(r)$ .


# Divergence of geodesics in Euclidean space

In Euclidean space, all pairs of geodesics diverge linearly.



# Divergence of geodesics in hyperbolic space

In hyperbolic space, all pairs of geodesics diverge exponentially.



# Divergence of geodesics in symmetric spaces

## Examples

1. In Euclidean space, all pairs of geodesics diverge linearly.
2. In hyperbolic space, all pairs of geodesics diverge exponentially.

# Divergence of geodesics in symmetric spaces

## Examples

1. In Euclidean space, all pairs of geodesics diverge linearly.
2. In hyperbolic space, all pairs of geodesics diverge exponentially.

## Theorem (Gromov)

*Let  $X$  be a symmetric space of noncompact type e.g.  $SL_n(\mathbb{R})/SO_n(\mathbb{R})$ . Then for all pairs of geodesics  $\gamma_1, \gamma_2$  with common basepoint, the function  $r \mapsto \text{div}(\gamma_1, \gamma_2, r)$  is either linear or exponential.*

# Divergence of geodesics in symmetric spaces

## Examples

1. In Euclidean space, all pairs of geodesics diverge linearly.
2. In hyperbolic space, all pairs of geodesics diverge exponentially.

## Theorem (Gromov)

*Let  $X$  be a symmetric space of noncompact type e.g.  $SL_n(\mathbb{R})/SO_n(\mathbb{R})$ . Then for all pairs of geodesics  $\gamma_1, \gamma_2$  with common basepoint, the function  $r \mapsto \text{div}(\gamma_1, \gamma_2, r)$  is either linear or exponential.*

Gromov suggested that the same dichotomy should hold in all CAT(0) spaces.

# Divergence of geodesics in symmetric spaces

## Examples

1. In Euclidean space, all pairs of geodesics diverge linearly.
2. In hyperbolic space, all pairs of geodesics diverge exponentially.

## Theorem (Gromov)

*Let  $X$  be a symmetric space of noncompact type e.g.  $SL_n(\mathbb{R})/SO_n(\mathbb{R})$ . Then for all pairs of geodesics  $\gamma_1, \gamma_2$  with common basepoint, the function  $r \mapsto \text{div}(\gamma_1, \gamma_2, r)$  is either linear or exponential.*

Gromov suggested that the same dichotomy should hold in all CAT(0) spaces.

It doesn't.



## Divergence for finitely generated groups

$G$  finitely generated group with finite generating set  $S$

## Divergence for finitely generated groups

$G$  finitely generated group with finite generating set  $S$

$X$  the Cayley graph of  $G$  w.r.t.  $S$ .

## Divergence for finitely generated groups

$G$  finitely generated group with finite generating set  $S$

$X$  the Cayley graph of  $G$  w.r.t.  $S$ . Assume  $X$  is one-ended.

## Divergence for finitely generated groups

$G$  finitely generated group with finite generating set  $S$   
 $X$  the Cayley graph of  $G$  w.r.t.  $S$ . Assume  $X$  is one-ended.

### Definition (Gersten 1994)

The **divergence** of  $G$  is the function

$$\operatorname{div}_G(r) := \sup_{x,y} \left( \inf_p \operatorname{length}(p) \right)$$

where

- ▶ the sup is over all pairs of points  $x, y \in X$  at distance  $r$  from  $e$
- ▶ the inf is over all paths  $p$  from  $x$  to  $y$  in  $X \setminus \operatorname{Ball}(e, r)$ .

## Divergence for finitely generated groups

$G$  finitely generated group with finite generating set  $S$   
 $X$  the Cayley graph of  $G$  w.r.t.  $S$ . Assume  $X$  is one-ended.

Definition (Gersten 1994)

The **divergence** of  $G$  is the function

$$\mathrm{div}_G(r) := \sup_{x,y} \left( \inf_p \mathrm{length}(p) \right)$$

where

- ▶ the sup is over all pairs of points  $x, y \in X$  at distance  $r$  from  $e$
- ▶ the inf is over all paths  $p$  from  $x$  to  $y$  in  $X \setminus \mathrm{Ball}(e, r)$ .

$G$  has **linear divergence** if  $\mathrm{div}_G(r) \simeq r$ , **quadratic divergence** if  $\mathrm{div}_G(r) \simeq r^2$ , etc, where

$$f \preceq g \iff \exists C > 0 \text{ s.t. } f(r) \leq Cg(Cr + C) + Cr + C$$

These rates of divergence are quasi-isometry invariants (Gersten).

## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

- ▶ quadratic divergence for certain free-by-cyclic groups [Gersten 1994]

## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

- ▶ quadratic divergence for certain free-by-cyclic groups [Gersten 1994]
- ▶ (geometric) 3-manifold groups have divergence either linear, quadratic or exponential; quadratic  $\iff$  graph manifold, exponential  $\iff$  hyperbolic piece [Gersten 1994, Kapovich–Leeb 1998]



## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

- ▶ quadratic divergence for certain free-by-cyclic groups [Gersten 1994]
- ▶ (geometric) 3-manifold groups have divergence either linear, quadratic or exponential; quadratic  $\iff$  graph manifold, exponential  $\iff$  hyperbolic piece [Gersten 1994, Kapovich–Leeb 1998]
- ▶ mapping class groups and Teichmüller space have quadratic divergence [Duchin–Rafi 2009]

## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

- ▶ quadratic divergence for certain free-by-cyclic groups [Gersten 1994]
- ▶ (geometric) 3-manifold groups have divergence either linear, quadratic or exponential; quadratic  $\iff$  graph manifold, exponential  $\iff$  hyperbolic piece [Gersten 1994, Kapovich–Leeb 1998]
- ▶ mapping class groups and Teichmüller space have quadratic divergence [Duchin–Rafi 2009]
- ▶ lattices in higher rank semisimple Lie groups conjectured to have linear divergence; proved in some cases e.g.  $SL(n, \mathbb{Z})$  [Drutu–Mozes–Sapir 2010]

## Previous results on divergence

Contrary to Gromov's expectation of linear or exponential divergence only:

- ▶ quadratic divergence for certain free-by-cyclic groups [Gersten 1994]
- ▶ (geometric) 3-manifold groups have divergence either linear, quadratic or exponential; quadratic  $\iff$  graph manifold, exponential  $\iff$  hyperbolic piece [Gersten 1994, Kapovich–Leeb 1998]
- ▶ mapping class groups and Teichmüller space have quadratic divergence [Duchin–Rafi 2009]
- ▶ lattices in higher rank semisimple Lie groups conjectured to have linear divergence; proved in some cases e.g.  $SL(n, \mathbb{Z})$  [Drutu–Mozes–Sapir 2010]
- ▶ right-angled Artin groups have divergence linear, quadratic or exponential [Abrams–Brady–Dani–Duchin–Young 2010, Behrstock–Charney 2012]

## Previous results on divergence

Constructions of finitely presented CAT(0) groups with divergence:

- ▶  $r^d$  for all integers  $d \geq 1$  [Macura 2011, Behrstock–Drutu 2011]

## Previous results on divergence

Constructions of finitely presented CAT(0) groups with divergence:

- ▶  $r^d$  for all integers  $d \geq 1$  [Macura 2011, Behrstock–Drutu 2011]
- ▶  $r^\alpha$  for  $\alpha$  irrational and dense in  $[2, \infty)$  [Brady–Tran 2020]

## Previous results on divergence

Constructions of finitely presented CAT(0) groups with divergence:

- ▶  $r^d$  for all integers  $d \geq 1$  [Macura 2011, Behrstock–Drutu 2011]
- ▶  $r^\alpha$  for  $\alpha$  irrational and dense in  $[2, \infty)$  [Brady–Tran 2020]
- ▶  $r^n \log r$  for all integers  $n \geq 1$  [Brady–Tran 2020]

# Right-angled Coxeter groups

$\Gamma$  finite simplicial graph with vertex set  $S$

# Right-angled Coxeter groups

$\Gamma$  finite simplicial graph with vertex set  $S$

The **right-angled Coxeter group** (RACG) associated to  $\Gamma$  is

$$W_\Gamma = \langle S \mid s^2 = 1 \forall s \in S \text{ and } (st)^2 = 1 \iff s \text{ and } t \text{ are adjacent in } \Gamma \rangle$$



# Right-angled Coxeter groups

$\Gamma$  finite simplicial graph with vertex set  $S$

The **right-angled Coxeter group** (RACG) associated to  $\Gamma$  is

$$W_\Gamma = \langle S \mid s^2 = 1 \forall s \in S \text{ and } (st)^2 = 1 \iff s \text{ and } t \text{ are adjacent in } \Gamma \rangle$$

equivalently

$$W_\Gamma = \langle S \mid s^2 = 1 \forall s \in S \text{ and } st = ts \iff s \text{ and } t \text{ are adjacent in } \Gamma \rangle$$

# Divergence in right-angled Coxeter groups

We consider  $W_\Gamma$  such that

- ▶  $\Gamma$  is triangle-free (simplifying assumption)

# Divergence in right-angled Coxeter groups

We consider  $W_\Gamma$  such that

- ▶  $\Gamma$  is triangle-free (simplifying assumption)
- ▶  $W_\Gamma$  is one-ended  $\iff \Gamma$  has no separating vertices or edges (special case of theorem of Davis)

# Divergence in right-angled Coxeter groups

We consider  $W_\Gamma$  such that

- ▶  $\Gamma$  is triangle-free (simplifying assumption)
- ▶  $W_\Gamma$  is one-ended  $\iff \Gamma$  has no separating vertices or edges (special case of theorem of Davis)


## Theorem 1 (Dani–T 2015)

1.  $W_\Gamma$  has linear divergence  $\iff \Gamma$  a join  $\iff W_\Gamma$  reducible.
2.  $W_\Gamma$  has quadratic divergence if and only if  $\Gamma$  is CFS and is not a join.

# Divergence in right-angled Coxeter groups

## Theorem 2 (Dani–T 2015)

*For all  $d \geq 1$ , the group  $W_{\Gamma_d}$  has divergence  $r^d$ .*



## Subsequent results for RACGs

Behrstock, Falgas-Ravry, Hagen and Susse generalised our  $\mathcal{CFS}$  condition to all  $\Gamma$ .

Theorem (Behrstock–Caprace–Hagen–Sisto, Sisto, Levcovitz)

1. *The divergence of  $W_\Gamma$  is either exponential ( $\iff$  the group is relatively hyperbolic) or bounded above by a polynomial ( $\iff$  the group is thick).*
2.  *$W_\Gamma$  has linear divergence  $\iff \Gamma$  is a join.*
3.  *$W_\Gamma$  has quadratic divergence  $\iff \Gamma$  is  $\mathcal{CFS}$ .*

## Subsequent results for RACGs

Behrstock, Falgas-Ravry, Hagen and Susse generalised our  $\mathcal{CFS}$  condition to all  $\Gamma$ .

Theorem (Behrstock–Caprace–Hagen–Sisto, Sisto, Levcovitz)

1. *The divergence of  $W_\Gamma$  is either exponential ( $\iff$  the group is relatively hyperbolic) or bounded above by a polynomial ( $\iff$  the group is thick).*
2.  *$W_\Gamma$  has linear divergence  $\iff \Gamma$  is a join.*
3.  *$W_\Gamma$  has quadratic divergence  $\iff \Gamma$  is  $\mathcal{CFS}$ .*

Levcovitz (2018) introduced the **hypergraph index** for a RACG.

## Subsequent results for RACGs

Behrstock, Falgas-Ravry, Hagen and Susse generalised our  $\mathcal{CFS}$  condition to all  $\Gamma$ .

Theorem (Behrstock–Caprace–Hagen–Sisto, Sisto, Levcovitz)

1. *The divergence of  $W_\Gamma$  is either exponential ( $\iff$  the group is relatively hyperbolic) or bounded above by a polynomial ( $\iff$  the group is thick).*
2.  *$W_\Gamma$  has linear divergence  $\iff \Gamma$  is a join.*
3.  *$W_\Gamma$  has quadratic divergence  $\iff \Gamma$  is  $\mathcal{CFS}$ .*

Levcovitz (2018) introduced the **hypergraph index** for a RACG.

Theorem (Levcovitz 2020)

*$W_\Gamma$  has hypergraph index  $h \iff W_\Gamma$  has divergence  $r^{h+1}$ .*



## Divergence for general Coxeter groups

$(W, S)$  Coxeter system such that  $W$  is one-ended

# Divergence for general Coxeter groups

$(W, S)$  Coxeter system such that  $W$  is one-ended

Theorem 3 (Dani–Naqvi–Soroko–T 2021)

$W$  has linear divergence  $\iff (W, S) = (W_1, S_1) \times (W_2, S_2)$   
where either:

1. both  $W_1$  and  $W_2$  are infinite, or
2.  $W_1$  is finite and  $(W_2, S_2)$  is irreducible affine of rank  $\geq 3$

# Divergence for general Coxeter groups

$(W, S)$  Coxeter system such that  $W$  is one-ended

**Theorem 3 (Dani–Naqvi–Soroko–T 2021)**

$W$  has linear divergence  $\iff (W, S) = (W_1, S_1) \times (W_2, S_2)$   
where either:

1. both  $W_1$  and  $W_2$  are infinite, or
2.  $W_1$  is finite and  $(W_2, S_2)$  is irreducible affine of rank  $\geq 3$

In all other cases,  $\text{div } W \asymp r^2$ .

# Divergence for general Coxeter groups

$(W, S)$  Coxeter system such that  $W$  is one-ended

**Theorem 3 (Dani–Naqvi–Soroko–T 2021)**

$W$  has linear divergence  $\iff (W, S) = (W_1, S_1) \times (W_2, S_2)$   
where either:

1. both  $W_1$  and  $W_2$  are infinite, or
2.  $W_1$  is finite and  $(W_2, S_2)$  is irreducible affine of rank  $\geq 3$

In all other cases,  $\text{div } W \succeq r^2$ .

Proof ingredients:

- ▶ General result of Kapovich–Leeb for locally compact CAT(0) spaces
- ▶ For  $W$  other than in (1) and (2), and  $w \in W$  a Coxeter element,  $w^\infty$  is a rank one geodesic (Caprace–Fujiwara)

# Hypergraph index for general Coxeter systems

# Hypergraph index for general Coxeter systems

Inductively define subsets of  $\mathcal{P}(S)$ :

# Hypergraph index for general Coxeter systems

Inductively define subsets of  $\mathcal{P}(S)$ :

$$\begin{aligned}\Lambda_0 = & \{T \subseteq S \mid W_T \text{ has linear divergence}\} \cup \\ & \{A \times K \mid W_A \text{ minimal nonspherical, } W_K \text{ spherical}\}\end{aligned}$$

For  $i \geq 0$

- ▶ for  $T, T' \in \Lambda_i$ , define  $T \sim T'$  if  $\exists$

$$T = T_0, T_1, \dots, T_m = T' \subset \Lambda_i$$

such that  $W_{T_j \cap T_{j+1}}$  is nonspherical

# Hypergraph index for general Coxeter systems

Inductively define subsets of  $\mathcal{P}(S)$ :

$$\Lambda_0 = \{T \subseteq S \mid W_T \text{ has linear divergence}\} \cup \\ \{A \times K \mid W_A \text{ minimal nonspherical, } W_K \text{ spherical}\}$$

For  $i \geq 0$

- ▶ for  $T, T' \in \Lambda_i$ , define  $T \sim T'$  if  $\exists$

$$T = T_0, T_1, \dots, T_m = T' \subset \Lambda_i$$

such that  $W_{T_j \cap T_{j+1}}$  is nonspherical

- ▶  $\Lambda_{i+1}$  is union of  $\sim$ -equivalence classes in  $\Lambda_i$ .



# Hypergraph index for general Coxeter systems

Inductively define subsets of  $\mathcal{P}(S)$ :

$$\begin{aligned}\Lambda_0 = & \{T \subseteq S \mid W_T \text{ has linear divergence}\} \cup \\ & \{A \times K \mid W_A \text{ minimal nonspherical, } W_K \text{ spherical}\}\end{aligned}$$

For  $i \geq 0$

- ▶ for  $T, T' \in \Lambda_i$ , define  $T \sim T'$  if  $\exists$

$$T = T_0, T_1, \dots, T_m = T' \subset \Lambda_i$$

such that  $W_{T_j \cap T_{j+1}}$  is nonspherical

- ▶  $\Lambda_{i+1}$  is union of  $\sim$ -equivalence classes in  $\Lambda_i$ .

Define **hypergraph index**  $h(W, S) = i$  if  $S \in \Lambda_i \setminus \Lambda_{i-1}$ , else  $h(W, S) = \infty$ .

# Hypergraph index and divergence

Theorem 4 (Dani–Naqvi–Soroko–T 2021)

# Hypergraph index and divergence

Theorem 4 (Dani–Naqvi–Soroko–T 2021)

1.  $h = \infty \iff W$  has exponential divergence  $\iff W$  is relatively hyperbolic

# Hypergraph index and divergence

## Theorem 4 (Dani–Naqvi–Soroko–T 2021)

1.  $h = \infty \iff W$  has exponential divergence  $\iff W$  is relatively hyperbolic
2.  $h = 0 \iff W$  has linear divergence

# Hypergraph index and divergence

## Theorem 4 (Dani–Naqvi–Soroko–T 2021)

1.  $h = \infty \iff W$  has exponential divergence  $\iff W$  is relatively hyperbolic
2.  $h = 0 \iff W$  has linear divergence
3. If  $h = 1$  then  $W$  has quadratic divergence

# Hypergraph index and divergence

## Theorem 4 (Dani–Naqvi–Soroko–T 2021)

1.  $h = \infty \iff W$  has exponential divergence  $\iff W$  is relatively hyperbolic
2.  $h = 0 \iff W$  has linear divergence
3. If  $h = 1$  then  $W$  has quadratic divergence
4. If  $1 \leq h < \infty$ , then  $W$  has divergence bounded above by  $r^{h+1}$ .

# Hypergraph index and divergence

## Theorem 4 (Dani–Naqvi–Soroko–T 2021)

1.  $h = \infty \iff W$  has exponential divergence  $\iff W$  is relatively hyperbolic
2.  $h = 0 \iff W$  has linear divergence
3. If  $h = 1$  then  $W$  has quadratic divergence
4. If  $1 \leq h < \infty$ , then  $W$  has divergence bounded above by  $r^{h+1}$ .

## Conjecture

If  $1 \leq h < \infty$  then  $W$  has divergence bounded below by  $r^{h+1}$ .

# Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.



## Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.

- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element

# Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.

- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element
- ▶ Looking at  $r$ -avoidant paths from  $\gamma_w(r)$  to  $\gamma_w(-r)$

# Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.

- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element
- ▶ Looking at  $r$ -avoidant paths from  $\gamma_w(r)$  to  $\gamma_w(-r)$
- ▶ A **wall** is the fixed set of a reflection

# Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.

- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element
- ▶ Looking at  $r$ -avoidant paths from  $\gamma_w(r)$  to  $\gamma_w(-r)$
- ▶ A **wall** is the fixed set of a reflection
- ▶ Estimate length of paths by counting “parallel” walls crossed

# Lower bound for divergence

Joint with Ignat Soroko, Dibyendu Roy.



- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element
- ▶ Looking at  $r$ -avoidant paths from  $\gamma_w(r)$  to  $\gamma_w(-r)$
- ▶ A **wall** is the fixed set of a reflection
- ▶ Estimate length of paths by counting “parallel” walls crossed
- ▶ First step: find “parallel” walls crossed by  $\gamma_w$  itself

# Lower bound for divergence


Joint with Ignat Soroko, Dibyendu Roy.

- ▶ Candidate geodesic:  $\gamma_w = w^\infty$  for  $w$  a Coxeter element
- ▶ Looking at  $r$ -avoidant paths from  $\gamma_w(r)$  to  $\gamma_w(-r)$
- ▶ A **wall** is the fixed set of a reflection
- ▶ Estimate length of paths by counting “parallel” walls crossed
- ▶ First step: find “parallel” walls crossed by  $\gamma_w$  itself
- ▶ Use root system to determine intersections of walls

## Computation in type $\tilde{A}_2$ ( $h = 0$ )





Computations for  $h = 1$








## Computations for $h = 1$



## Computations for $h = 1$



## Computations for $h = 1$



Thank you for your attention.